

Ticagrelor–Aspirin Versus Clopidogrel–Aspirin Among *CYP2C19* Loss-of-Function Carriers With Minor Stroke or Transient Ischemic Attack in Relation to Renal Function: A Post Hoc Analysis of the CHANCE-2 Trial

Anxin Wang, PhD*; Xuewei Xie, MD, PhD*; Xue Tian, PhD; S. Claiborne Johnston, MD, PhD; Hao Li, MD, PhD; Philip M. Bath, DSc; Yingting Zuo, PhD; Jing Jing, MD, PhD; Jinxi Lin, MD, PhD; Yilong Wang, MD, PhD; Xingquan Zhao, MD, PhD; Zixiao Li, MD, PhD; Yong Jiang, MD, PhD; Liping Liu, MD, PhD; Xia Meng, MD; and Yongjun Wang, MD

Background: Evidence on the risk–benefit ratio of dual antiplatelet therapies among patients with stroke and impaired renal function is limited and inconsistent.

Objective: To investigate the effect of renal function on the efficacy and safety of ticagrelor–aspirin versus clopidogrel–aspirin treatment.

Design: Post hoc analysis of a multicenter, randomized, double-blind, placebo-controlled trial. (ClinicalTrials.gov: NCT04078737)

Setting: 202 centers in China.

Patients: *CYP2C19* loss-of-function allele carriers with minor stroke or transient ischemic attack.

Intervention: Ticagrelor–aspirin and clopidogrel–aspirin.

Measurements: Renal function was evaluated by estimated glomerular filtration rate (eGFR) levels. The primary efficacy and safety outcomes were recurrent stroke and severe or moderate bleeding within 90 days, respectively.

Results: Among 6378 patients, 4050 (63.5%) had normal (eGFR ≥ 90 mL/min/1.73 m²), 2010 (31.5%) had mildly decreased (eGFR 60 to 89 mL/min/1.73 m²), and 318 (5.0%) had moderately to severely decreased (eGFR <60 mL/min/1.73 m²) renal function.

The corresponding differences in recurrent stroke between ticagrelor–aspirin and clopidogrel–aspirin for normal, mildly decreased, and moderately to severely decreased renal function was –2.8 percentage points (95% CI, –4.4 to –1.3 percentage points) (hazard ratio [HR], 0.63 [CI, 0.49 to 0.81]), –0.2 percentage point (CI, –2.4 to 2.0 percentage points) (HR, 0.98 [CI, 0.69 to 1.39]), and 3.7 percentage points (CI, –2.3 to 10.1 percentage points) (HR, 1.31 [CI, 0.48 to 3.55]), respectively. Rates of severe or moderate bleeding did not substantially differ by treatment assignments across eGFR categories.

Limitation: Renal function was only evaluated by using eGFR, and the proportion of patients with severely decreased renal function was low.

Conclusion: Patients with normal, rather than impaired, renal function received greater benefit from ticagrelor–aspirin versus clopidogrel–aspirin.

Primary Funding Source: Ministry of Science and Technology of the People's Republic of China.

Ann Intern Med. doi:10.7326/M22-1667

Annals.org

For author, article, and disclosure information, see end of text.

This article was published at Annals.org on 1 November 2022.

* Drs. A. Wang and Xie contributed equally to this work.

Impaired renal function is associated with abnormalities in platelet function, which may explain increases in both thrombotic and hemorrhagic complications in patients with stroke (1–5). Antiplatelet therapies can reduce thrombotic risk in patients with impaired renal function but come at the expense of impaired hemostasis. This may alter the risk–benefit ratio with antiplatelet therapies in patient with stroke and impaired renal function. Therefore, determining the optimal antiplatelet strategies in this population is of utmost importance.

Dual antiplatelet therapy with clopidogrel–aspirin is often recommended for preventing stroke (6–8). Ticagrelor, a reversible and direct-acting oral antagonist of the P2Y₁₂ inhibitor, can provide greater, faster, and more consistent P2Y₁₂ inhibition than clopidogrel (9, 10). Ticagrelor

has been shown to be an effective antiplatelet therapy for the prevention of recurrent stroke (11), particularly in those carrying *CYP2C19* loss-of-function (LOF) alleles (12, 13). Reduced renal clearance of clopidogrel (and less so ticagrelor) could increase the risk for increased plasma concentrations in patients with impaired renal function, so renal function needs to be considered when selecting optimal antiplatelet therapy. Some studies suggested that patients with impaired renal function may not derive the same degree of benefit from clopidogrel therapy as those with normal renal function (14, 15). In contrast, some studies have suggested that patients with impaired renal function received more benefit from clopidogrel or ticagrelor (16, 17). In addition, uncertainties remain about whether the benefit of ticagrelor–aspirin versus clopidogrel–aspirin is in relation to renal function among *CYP2C19* LOF allele carriers with minor ischemic stroke or transient ischemic attack (TIA).

Using data from the CHANCE-2 (Clopidogrel in High-Risk Patients with Acute Nondisabling Cerebrovascular Events-II) trial, we investigated the efficacy and safety of

See also:

Web-Only
Supplement

ticagrelor-aspirin versus clopidogrel-aspirin in patients with minor stroke or TIA who carried *CYP2C19* LOF alleles with different renal function evaluated by estimated glomerular filtration rate (eGFR) levels.

METHODS

Study Design and Populations

This study is a post hoc analysis of the CHANCE-2 trial. Details on the design, protocol, and primary results of CHANCE-2 have been published elsewhere (12, 18) (the study protocol, and the statistical analysis plan, can also be found at [Annals.org](https://www.annals.org)). Briefly, CHANCE-2 was a randomized, double-blind, controlled trial conducted at 202 centers across mainland China from 23 September 2019 to 22 March 2021 (ClinicalTrials.gov: NCT04078737). A total of 6412 patients who met the following inclusion criteria were enrolled; they: (1) were aged 40 years or older; (2) had a mild acute ischemic stroke (National Institutes of Health Stroke Score of ≤ 3) or a high-risk TIA (ABCD² [age, blood pressure, clinical features, duration of TIA, and presence or absence of diabetes] score of ≥ 4); (3) were carriers of *CYP2C19* LOF alleles; (4) had the trial drug administered within 24 hours of symptom onset; and (5) signed informed consent. The protocol of the trial was approved by the ethics committee at Beijing Tiantan Hospital (institutional review board approval number: KY2019-035-02) and each participating site. All participants or their representatives provided written informed consent before enrollment.

Randomization and Treatment

Within 24 hours after symptom onset, eligible patients carrying *CYP2C19* LOF alleles were randomly assigned in a 1:1 ratio to receive ticagrelor-aspirin or clopidogrel-aspirin. Patients were randomly assigned a number corresponding to a medication kit that was given to each patient. Patients in the ticagrelor-aspirin group received the clopidogrel placebo and a 180-mg loading dose of ticagrelor on day 1 followed by 90 mg twice daily for days 2 to 90. Patients in the clopidogrel-aspirin group received the ticagrelor placebo and a 300-mg loading dose of clopidogrel, followed by 75 mg daily together for days 2 to 90. Both groups received a 75- to 300-mg loading dose of aspirin on day 1, followed by 75 mg daily for 21 days.

Calculation of eGFR

Venous blood samples were obtained before randomization and were sent for laboratory analysis of creatinine concentration. The eGFR was calculated using the Chronic Kidney Disease Epidemiology Collaboration creatinine equation (CKD-EPI) (19): $eGFR = 141 \times \min(SCr/k, 1)^\alpha \times \max(SCr/k, 1)^{-1.209} \times 0.993^{Age} \times 1.018$ (if the patient is female), where *SCr* is serum creatinine, *k* is 0.7 for female patients and 0.9 for male patients, α is -0.329 for female patients and -0.411 for male patients, *min* is the minimum of *SCr*/*k* or 1, and *max* indicates the maximum of *SCr*/*k* or 1. The CKD-EPI China equation was calculated with a coefficient of 1.1 (20). According to the National Kidney Foundation Kidney Disease Outcomes Quality

Initiative (NKF-KDOQI) guidelines (21, 22), normal renal function was defined as an eGFR of at least 90 mL/min/1.73 m², mildly decreased renal function was defined as an eGFR of 60 to 89 mL/min/1.73 m², moderately decreased renal function was defined as an eGFR of 30 to 59 mL/min/1.73 m², and severely decreased renal function was defined as an eGFR of less than 30 mL/min/1.73 m².


Outcomes Assessment

The primary outcome was a new ischemic or hemorrhagic stroke within 90 days. Secondary outcomes included new stroke within 30 days, composite vascular events (stroke, TIA, myocardial infarction, and vascular death), ischemic stroke, disabling stroke (with a subsequent modified Rankin Scale [mRS] score of 2 or higher; range 0 to 6, with higher scores reflecting greater handicap) at day 90, and ordinal severity of stroke or TIA through 90 days of follow-up. For ordinal severity of stroke or TIA, severity was measured using a 6-level ordered categorical scale that incorporates subsequent stroke or TIA events and mRS score at day 90 (5): fatal stroke (stroke with subsequent mRS score of 6), severe stroke (stroke with subsequent mRS score of 4 or 5), moderate stroke (stroke with subsequent mRS score of 2 or 3), mild stroke (stroke with subsequent mRS score of 0 or 1), TIA, and no stroke or TIA.

The primary safety outcome was severe or moderate bleeding defined by the GUSTO (Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Coronary Arteries) criteria within 90 days (23). Secondary safety outcomes included any bleeding, death, adverse events, and severe adverse events through 90 days of follow-up.

Statistical Analysis

Continuous variables are presented as median with interquartile range, and categorical variables as frequencies and percentages. The differences in the proportions for the dichotomous outcomes between treatment groups, and their corresponding 95% CIs, were estimated based on Newcombe-Wilson (24), with stratification by eGFR category. Kaplan-Meier analysis was used to calculate the cumulative incidence of the primary outcome during 90 days of follow-up for each eGFR category. Differences in the outcome end points during the 90-day follow-up period were assessed using a Cox proportional hazards regression model, with study centers set as a random effect, and hazard ratios (HRs) with 95% CIs reported. When there were several events of the same type, the time to the first event was used in the model. Patients without any events during 90-day follow-up were censored at the time of termination of the trial or nonvascular death. Similar methods were used for the comparison of the secondary outcomes of new stroke events, clinical vascular events, ischemic stroke, and disabling stroke, and for comparison of the safety outcomes. Shift analysis was performed for the secondary outcome of ordinal stroke or TIA between the 2 treatment groups using ordinal logistic regression, and the common odds ratio and 95% CI reported. To test the robustness of the findings, sensitivity analyses were performed by calculating the eGFR using the CKD-EPI for

Figure 1. The flow chart of the study.

ABCD² = age, blood pressure, clinical features, duration of TIA, and presence or absence of diabetes; eGFR = estimated glomerular filtration rate; NIHSS = National Institutes of Health Stroke Scale; TIA = transient ischemic attack.

the Chinese population and in the per protocol population. All statistical analyses were performed with SAS statistical software, version 9.4 (SAS Institute).

Role of the Funding Source

The work was supported by the Ministry of Science and Technology of the People's Republic of China, the Beijing Municipal Science and Technology Commission, the Chinese Stroke Association, the National Science and Technology Major Project, and the Beijing Municipal Administration of Hospitals Incubating Program. The funders did not influence study design, conduct, or reporting.

RESULTS

Baseline Characteristics

Of the 6412 eligible patients recruited to the CHANCE-2 trial, 6378 (99.5%) with eGFR measurement were analyzed in the current study (Figure 1). The median age of enrolled patients was 64.5 years (interquartile range, 57.0 to 71.4 years), and 2165 (33.9%) were women. Overall, 4050 patients (63.5%) had normal renal function (eGFR ≥ 90 mL/min/1.73 m²), 2010 (31.5%) had mildly decreased renal function (eGFR of 60 to 89 mL/min/1.73 m²), 309 (4.8%) had moderately decreased renal function (eGFR of 30 to 59 mL/min/1.73 m²), and 9 (0.1%) had severely decreased renal function (eGFR <30 mL/min/1.73 m²). Considering the small sample size of patients with moderately and severely decreased renal

function, those patients were combined into 1 group as moderately to severely decreased renal function (eGFR <60 mL/min/1.73 m²). The baseline characteristics in the ticagrelor–aspirin and clopidogrel–aspirin groups across the 3 eGFR categories were well balanced (Table 1).

Efficacy Outcomes

The primary efficacy outcome of recurrent stroke within 90 days occurred in 189 patients (5.9%) receiving ticagrelor–aspirin and 243 patients (7.6%) receiving clopidogrel–aspirin. Ticagrelor–aspirin compared with clopidogrel–aspirin was associated with a reduced rate of recurrent stroke in patients with normal renal function (5.2% vs. 8.1%; difference, –2.8 percentage points [95% CI, –4.4 to –1.3 percentage points]; HR, 0.63 [CI, 0.49 to 0.81]), but not in those with mildly decreased renal function (6.7% vs. 6.9%; difference, –0.2 percentage points [CI, –2.4 to 2.0 percentage points]; HR, 0.98 [CI, 0.69 to 1.39]), or those with moderately to severely decreased renal function (9.8% vs. 6.1%; difference, 3.7 percentage points [CI, –2.3 to 10.1 percentage points]; HR, 1.31 [CI, 0.48 to 3.55]) (Table 2 and Figure 2). Similar results were present for the secondary outcomes of combined vascular event, ischemic stroke, and ordinal stroke or TIA within 90 days of follow-up (Table 2).

Results of the sensitivity analysis by calculating the eGFR using the CKD-EPI for the Chinese population were consistent with the primary analysis, showing that the difference in the rate of recurrent stroke between the

Table 1. Baseline Characteristics According to eGFR Category

Characteristics	eGFR <60 mL/min/ 1.73 m ² (N = 318)		eGFR 60-89 mL/min/ 1.73 m ² (N = 2010)		eGFR ≥90 mL/min/ 1.73 m ² (N = 4050)	
	T-A (N = 153)	C-A (N = 165)	T-A (N = 1028)	C-A (N = 982)	T-A (N = 2006)	C-A (N = 2044)
Median age (IQR), y	75.3 (67.2-80.3)	72.8 (65.1-80.5)	71.5 (64.5-77.3)	71.1 (65.3-76.9)	61.5 (54.8-67.1)	61.1 (54.7-66.6)
Female sex, n (%)	64 (41.8)	68 (41.2)	377 (36.7)	372 (37.9)	645 (32.2)	639 (31.3)
Han ethnicity, n (%)	151 (98.7)	159 (96.4)	1008 (98.1)	959 (97.7)	1967 (98.1)	2004 (98.0)
Body mass index, kg/m ²	23.9 (22.1-25.7)	23.5 (22.2-25.7)	24.4 (22.5-26.6)	24.2 (22.3-26.1)	24.6 (22.9-26.7)	24.5 (22.8-26.6)
Median blood pressure (IQR), mm Hg						
Systolic	148.0 (135.0-173.0)	150.0 (140.0-164.5)	148.0 (136.0-161.5)	148.0 (135.0-162.5)	148.5 (136.0-162.0)	148.0 (136.0-160.0)
Diastolic	81.0 (75.0-93.5)	84.0 (80.0-92.0)	84.0 (78.0-92.0)	84.5 (78.0-93.0)	88.0 (80.0-96.0)	87.5 (80.0-96.0)
Medical history, n (%)						
Hypertension	125 (81.7)	139 (84.2)	776 (75.5)	760 (77.4)	1441 (71.8)	1465 (71.7)
Diabetes mellitus	64 (41.8)	65 (39.4)	319 (31.0)	260 (26.5)	646 (32.2)	678 (33.2)
Dyslipidemia	35 (22.9)	48 (29.1)	288 (28.0)	285 (29.0)	562 (28.0)	562 (27.5)
Previous ischemic stroke	35 (22.9)	43 (26.1)	237 (23.1)	224 (22.8)	395 (19.7)	409 (20.0)
Previous TIA	3 (2.0)	0 (0.0)	13 (1.3)	15 (1.5)	30 (1.5)	27 (1.3)
Myocardial infarction	3 (2.0)	3 (1.8)	19 (1.8)	14 (1.4)	31 (1.5)	25 (1.2)
Current smoking, n (%)	36 (23.5)	37 (22.4)	265 (25.8)	232 (23.6)	690 (34.4)	710 (34.7)
CYP2C19 LOF allele carriers, n (%)						
Intermediate metabolizers	126 (82.4)	136 (82.4)	799 (77.7)	765 (77.9)	1548 (77.2)	1601 (78.3)
Poor metabolizers	27 (17.6)	29 (17.6)	229 (22.3)	217 (22.1)	458 (22.8)	443 (21.7)
Qualifying event, n (%)						
Ischemic stroke	126 (82.4)	139 (84.2)	795 (77.3)	768 (78.2)	1645 (82.0)	1665 (81.5)
TIA	27 (17.6)	26 (15.8)	233 (22.7)	214 (21.8)	361 (18.0)	379 (18.5)
Median NIHSS score in patients with qualifying ischemic stroke (IQR)*	2 (1-2)	2 (1-3)	2 (1-3)	2 (1-3)	2 (1-3)	2 (1-3)
Median ABCD ² score in patients with qualifying TIA (IQR)†	5 (5-6)	5 (4-6)	5 (4-5)	5 (4-5)	4 (4-5)	4 (4-5)
Previous antiplatelet therapy, n (%)‡	23 (15.0)	20 (12.1)	144 (14.0)	113 (11.5)	217 (10.8)	226 (11.1)
Previous lipid-lowering therapy, n (%)‡	16 (10.5)	14 (8.5)	98 (9.5)	76 (7.7)	143 (7.1)	149 (7.3)

ABCD² = age, blood pressure, clinical features, duration of TIA, and presence or absence of diabetes; C-A = clopidogrel-aspirin; eGFR = estimated glomerular filtration rate; IQR = interquartile range; LOF = loss of function; NIHSS = National Institutes of Health Stroke Scale; T-A = ticagrelor-aspirin; TIA = transient ischemic attack.

* National Institutes of Health Stroke Scale scores range from 0 to 42, with higher scores indicating more severe stroke.

† ABCD² score assesses the risk for stroke on the basis of age, blood pressure, clinical features, duration of TIA, and presence or absence of diabetes, with scores ranging from 0 to 7 and higher scores indicating greater risk.

‡ Medication within 1 mo before symptom onset.

ticagrelor-aspirin and clopidogrel-aspirin groups was -2.4 percentage points [CI, -3.8 to -1.1 percentage points] (HR, 0.68 [CI, 0.54 to 0.84]) in patients with normal renal function (Supplement Table 1 and Supplement Figure 1, available at [Annals.org](https://annals.org)). In addition, the per protocol analysis yielded similar results to the intention-to-treat analysis; the HR for recurrent stroke in the ticagrelor-aspirin group compared with the clopidogrel-aspirin group was 0.61 (CI, 0.48 to 0.79) in patients with normal renal function (Supplement Table 2 and Supplement Figure 2, available at [Annals.org](https://annals.org)).

Safety Outcomes


The rate of primary safety outcome of severe or moderate bleeding in the ticagrelor-aspirin group and the clopidogrel-aspirin group was similar in patients with normal renal function (0.2% vs. 0.3%; difference, -0.1 percentage point [CI, -0.5 to 0.2 percentage point], HR, 0.59 [CI, 0.17 to 2.01]), mildly decreased renal function (0.4% vs. 0.3%; difference, 0.1 percentage point [CI, -0.6 to 0.7 percentage point], HR, 1.28 [CI, 0.28 to 5.75]), and

moderately to severely decreased renal function (0.7% vs. 0.6%; difference, 0.0 percentage point [CI, -2.8 to 3.0 percentage points]) (Table 3). Similar results were yielded for second safety outcomes. Sensitivity analyses were consistent with the main analysis (Supplement Tables 1 and 2).

DISCUSSION

Based on the CHANCE-2 trial, our study found that ticagrelor-aspirin, compared with clopidogrel-aspirin, substantially reduced the risk for recurrent stroke within 90 days of follow-up in patients with normal renal function, but this benefit was not apparent in those with mildly or moderately to severely decreased renal function. Meanwhile, there was no absolute increase in severe or moderate bleeding events with ticagrelor-aspirin treatment across eGFR categories although this was based on small numbers.

Many post hoc analyses have evaluated the effect of renal function on the efficacy and safety of antiplatelet therapies and yielded divergent results in this context. Some studies observed a substantial benefit of intensive

Figure 2. Cumulative probability of stroke according to treatment and eGFR category.

C-A = clopidogrel-aspirin; eGFR = estimated glomerular filtration rate; T-A = ticagrelor-aspirin.

antiplatelet therapies among patients with normal renal function. For example, in the CREDO (Clopidogrel for the Reduction of Events During Observation) trial, clopidogrel versus placebo reduced the composite of death, myocardial infarction, and stroke in patients with normal renal function and acute coronary syndrome after percutaneous coronary intervention, but with a trend in the opposite direction with an absolute increased event rate in patients with mild or moderate renal dysfunction (14). A post

hoc analysis of the CHANCE trial found that clopidogrel plus aspirin compared with aspirin alone in patients with normal renal function and mild renal insufficiency resulted in a substantial reduction in new stroke events and combined vascular events at 90 days of follow-up, but this benefit was not apparent in patients with moderate chronic kidney disease (CKD) (15). In accordance with these studies, our study was conducted among *CYP2C19* LOF allele carriers with minor ischemic stroke or TIA, showing that ticagrelor-aspirin, of which ticagrelor can provide more consistent P2Y₁₂ inhibition than clopidogrel (9, 10), was associated with a lower risk for recurrent stroke in patients with normal renal function compared with clopidogrel-aspirin, whereas the benefit was not observed in patients with mildly or moderately to severely decreased renal function.

However, as opposed to these results, some studies showed reduced or lack of effect with intensive antiplatelet therapies among patients with normal renal function. In the CURE (Clopidogrel in Unstable Angina to Prevent Recurrent Events) study, there was a modest absolute and relative reduction in the primary ischemic end point with clopidogrel versus placebo among patients with renal dysfunction compared with those with normal renal function, although without interaction (16). Subgroup analysis of the PLATO (Platelet Inhibition and Patient Outcomes) trial showed that ticagrelor significantly reduced the rate of ischemic end points and mortality compared with clopidogrel in patients with acute coronary syndrome with CKD; also, the interactions between creatinine clearance and randomized treatment on any of the outcome variables were nonsignificant (17).

Although the reasons for these inconsistent results are unclear, plausible explanations may include the highly heterogeneous target populations across these studies, as well as the different treatment assignments and trial design paradigms. In addition, potential mechanisms underlying our results may be a synergistic relationship between the thrombotic effects of renal dysfunction and the antithrombotic effects of dual antiplatelet treatment. First, decreased renal function is characterized as a state with a prothrombotic tendency, and is associated with anemia, homocysteinemia, reduced nitric oxide, oxidative stress, inflammation, and conditions promoting coagulation; all of these pathologic processes may be related to the development of recurrent stroke in the course of decreased renal function (25–27). The levels of platelet inhibition from different antiplatelet therapies may not be sufficient for adequate protection against ischemic events in these patients at high risk. Second, differences in the pharmacodynamic and pharmacokinetic profiles of ticagrelor and clopidogrel (28) mean the excretion of ticagrelor is less dependent on renal function compared with clopidogrel (29–31). One pharmacologic study showed that 26.5% of ticagrelor is excreted through the kidney, and the recovery of ticagrelor and its active metabolites in the urine is less than 1% of the dose (32). As for clopidogrel, almost 50% of clopidogrel as well as part of its active metabolite was excreted in the urine (33); hence, renal clearance is of minor importance in the excretion of ticagrelor compared with clopidogrel. For patients with decreased renal function, the excretion of clopidogrel and its active

Table 2. Efficacy Outcomes of Patients With Different Antiplatelet Therapies Stratified by eGFR Category

Outcome	eGFR <60 mL/min/1.73 m ²				eGFR 60-89 mL/min/1.73 m ²				eGFR ≥90 mL/min/1.73 m ²			
	T-A, %	C-A, %	HR/OR (95% CI)	RD (95% CI), %	T-A, %	C-A, %	HR/OR (95% CI)	RD (95% CI), %	T-A, %	C-A, %	HR/OR (95% CI)	RD (95% CI), %
Primary outcome												
Stroke	15 (9.8)	10 (6.1)	1.31 (0.48 to 3.55)	3.7 (-2.3 to 10.1)	69 (6.7)	68 (6.9)	0.98 (0.69 to 1.39)	-0.2 (-2.4 to 2.0)	105 (5.2)	165 (8.1)	0.63 (0.49 to 0.81)	-2.8 (-4.4 to -1.3)
Secondary outcome												
Stroke within 30 d	14 (9.2)	8 (4.8)	1.53 (0.54 to 4.32)	4.3 (-1.4 to 10.4)	51 (5.0)	58 (5.9)	0.85 (0.57 to 1.25)	-0.9 (-3.0 to 1.8)	89 (4.4)	139 (6.8)	0.64 (0.49 to 0.84)	-2.4 (-3.8 to -0.9)
Composite vascular events*	17 (11.1)	12 (7.3)	1.35 (0.53 to 3.47)	3.8 (-2.6 to 10.5)	83 (8.1)	85 (8.7)	0.97 (0.71 to 1.33)	-0.6 (-3.0 to 1.8)	126 (6.3)	195 (9.5)	0.65 (0.52 to 0.81)	-3.3 (-4.9 to -1.6)
Ischemic stroke	15 (9.8)	10 (6.1)	1.31 (0.48 to 3.55)	3.7 (-2.3 to 10.1)	67 (6.5)	68 (6.9)	0.95 (0.67 to 1.35)	-0.4 (-2.6 to 1.8)	105 (5.3)	160 (7.8)	0.65 (0.51 to 0.84)	-2.6 (-4.1 to -1.1)
Disabling stroke†	6 (3.9)	7 (4.2)	0.61 (0.11 to 3.31)	-0.3 (-5.1 to 4.6)	39 (3.8)	24 (2.4)	1.43 (0.85 to 2.40)	1.3 (-0.2 to 2.9)	51 (2.5)	61 (3.0)	0.82 (0.56 to 1.19)	-0.4 (-1.5 to 0.6)
Ordinal stroke or TIA‡	-	-	1.46 (0.67 to 3.20)	NA	-	-	1.00 (0.73 to 1.39)	NA	-	-	0.65 (0.51 to 0.82)	NA
Fatal stroke (mRS 6)	0 (0.0)	1 (0.6)	-	-	2 (0.2)	2 (0.2)	-	-	2 (0.1)	5 (0.2)	-	-
Severe stroke (mRS 4-5)	3 (2.0)	3 (1.8)	-	-	10 (1.0)	4 (0.4)	-	-	16 (0.8)	14 (0.7)	-	-
Moderate stroke (mRS 2-3)	3 (2.0)	3 (1.8)	-	-	27 (2.6)	18 (1.8)	-	-	33 (1.6)	42 (2.1)	-	-
Mild stroke (mRS 0-1)	9 (5.9)	3 (1.8)	-	-	30 (2.9)	44 (4.5)	-	-	54 (2.7)	104 (5.1)	-	-
TIA	1 (0.7)	2 (1.2)	-	-	12 (1.2)	10 (1.0)	-	-	20 (1.0)	27 (1.3)	-	-
No stroke or TIA	137 (89.5)	153 (92.7)	-	-	947 (92.1)	904 (92.1)	-	-	1881 (93.8)	1852 (90.6)	-	-

C-A = clopidogrel-aspirin; eGFR = estimated glomerular filtration rate; HR = hazard ratio; mRS = modified Rankin Scale; NA = not applicable; OR = odds ratio; RD = risk difference; T-A = ticagrelor-aspirin; TIA = transient ischemic attack.

* Composite vascular events include ischemic stroke, hemorrhagic stroke, TIA, myocardial infarction, vascular death.

† A stroke was defined as disabling if the patient had a subsequent score on the mRS of >1 (indicating death or any degree of disability).

‡ Severity measured using a 6-level ordered categorical scale that incorporates subsequent stroke or TIA events and mRS score at day 90. The common OR is shown rather than the HR.

metabolites will be limited, leading to an increase in half-life and higher peak concentration in the body. As a result, the benefit of ticagrelor over clopidogrel may not be observed in patients with decreased renal function.

For the safety outcomes of bleeding risk, a substudy of the PLATO study also found that major bleedings were not significantly increasing in the ticagrelor group compared with the clopidogrel group (17). The TWILIGHT-CKD (The Ticagrelor With Aspirin or Alone in High Risk Patients After Coronary Intervention) trial showed that among patients with CKD undergoing percutaneous coronary intervention, ticagrelor monotherapy reduced the risk for bleeding without a substantial increase in ischemic events compared with ticagrelor plus aspirin (34). In line with these studies, our study found that patients receiving ticagrelor-aspirin did not show an absolute increase in severe or moderate bleeding events across different eGFR categories, although the number of bleeding events was relatively small in our study. However, it should be noted that the incidence of total bleeding, mainly mild bleeding, was greater with ticagrelor-aspirin in different eGFR categories. In addition, several previous studies have demonstrated that patients with impaired renal function have a higher bleeding tendency regardless of antiplatelet therapies (35-37). Taken together, these findings indicated that the bleeding risk should be carefully assessed and monitored in clinical utility of antiplatelet therapies.

There were several limitations to the study. First, renal function was defined by eGFR only, with no data available

on the presence of albuminuria or proteinuria. Although it will be more precise to diagnose CKD based on the combination of eGFR and albuminuria-proteinuria, the collection and measurement of urine samples in the acute stage of stroke-TIA is challenging in a large population. Ideally, albuminuria and proteinuria would be assessed in future investigations. Second, only a minority of patients had moderately to severely decreased renal function, thus caution is needed when interpreting the efficacy and safety of dual antiplatelet therapy for patients with stroke with moderately to severely decreased renal function. However, although specific recommendations for antiplatelet therapy for this special population are not available, the present study may provide some valuable information. A prospective and well-designed study in patients with stroke with impaired renal function would be needed for further evaluation. Third, this study was a post hoc analysis, which increases the risk for a type I error, so our result needed to be confirmed by other studies (38). Finally, all patients in the CHANCE-2 trial were Chinese, which may limit the generalizability of the findings to other populations.

In conclusion, based on the CHANCE-2 trial, our study showed that among *CYP2C19* LOF carriers with minor stroke or TIA, ticagrelor-aspirin compared with clopidogrel-aspirin was associated with a reduced risk for recurrent stroke and without any substantial increase in severe or moderate bleeding events among patients with normal renal function, whereas patients with impaired renal function

Table 3. Safety Outcomes of Patients With Different Antiplatelet Therapies Stratified by eGFR Category

Outcome	eGFR <60 mL/min/1.73 m ²				eGFR 60-89 mL/min/1.73 m ²				eGFR ≥90 mL/min/1.73 m ²			
	T-A, %	C-A, %	HR (95% CI)	RD (95% CI), %	T-A, %	C-A, %	HR (95% CI)	RD (95% CI), %	T-A, %	C-A, %	HR (95% CI)	RD (95% CI), %
Primary safety outcome												
Severe or moderate bleeding*	1 (0.7)	1 (0.6)	NA	0.0 (−2.8 to 3.0)	4 (0.4)	3 (0.3)	1.28 (0.28 to 5.75)	0.1 (−0.6 to 0.7)	4 (0.2)	7 (0.3)	0.59 (0.17 to 2.01)	−0.1 (−0.5 to 0.2)
Fatal bleeding	0 (0.0)	0 (0.0)	NA	0.0 (−2.3 to 2.4)	1 (0.1)	1 (0.1)	0.77 (0.05 to 12.77)	0.0 (−0.5 to 0.5)	2 (0.1)	2 (0.1)	0.92 (0.13 to 6.57)	0.0 (−0.3 to 0.3)
Intracranial hemorrhage	0 (0.0)	0 (0.0)	NA	0.0 (−2.3 to 2.4)	2 (0.2)	1 (0.1)	1.72 (0.15 to 19.31)	0.1 (−0.4 to 0.6)	1 (0.0)	5 (0.2)	0.21 (0.02 to 1.78)	−0.2 (−0.5 to 0.1)
Secondary safety outcome												
Any bleeding	8 (5.3)	3 (1.8)	2.39 (0.41 to 13.88)	3.4 (−0.8 to 8.3)	49 (4.8)	29 (3.0)	1.46 (0.91 to 2.36)	1.8 (0.1 to 3.5)	113 (5.6)	48 (2.3)	2.49 (1.76 to 3.52)	3.3 (2.1 to 4.5)
Mild bleeding*	7 (4.6)	2 (1.2)	1.81 (0.28 to 11.76)	3.4 (−0.5 to 8.0)	45 (4.4)	26 (2.6)	1.49 (0.90 to 2.46)	1.7 (0.1 to 3.4)	109 (5.4)	41 (2.0)	2.85 (1.97 to 4.12)	3.4 (2.3 to 4.6)
Death	2 (1.3)	1 (0.6)	3.00 (0.25 to 35.79)	0.7 (−2.2 to 4.1)	3 (0.3)	11 (1.1)	0.31 (0.08 to 1.16)	−0.8 (−1.7 to −0.1)	4 (0.2)	6 (0.3)	0.66 (0.18 to 2.35)	−0.1 (−0.5 to 0.3)

C-A = clopidogrel-aspirin; eGFR = estimated glomerular filtration rate; HR = hazard ratio; NA = not applicable; RD = risk difference; T-A = ticagrelor-aspirin. * Severe or moderate bleeding and mild bleeding were defined according to GUSTO (Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Coronary Arteries) criteria.

did not derive the same benefit from ticagrelor-aspirin. The findings suggest that renal function should be considered when deciding on the use of ticagrelor-aspirin versus clopidogrel-aspirin.

From Department of Neurology, Beijing Tiantan Hospital, Capital Medical University, Beijing, and China National Clinical Research Center for Neurological Diseases, Beijing, China (A.W., X.X., H.L., J.J., J.L., Y.W., X.Z., Z.L., Y.J., L.L., X.M.); Department of Neurology, Beijing Tiantan Hospital, Capital Medical University, Beijing, China National Clinical Research Center for Neurological Diseases, Beijing, Department of Epidemiology and Health Statistics, School of Public Health, Capital Medical University, Beijing, and Beijing Municipal Key Laboratory of Clinical Epidemiology, Beijing, China (X.T., Y.Z.); Dell Medical School, The University of Texas at Austin, Austin, Texas (S.C.J.); Stroke Trials Unit, Mental Health & Clinical Neuroscience, University of Nottingham, Nottingham, United Kingdom (P.M.B.); and Department of Neurology, Beijing Tiantan Hospital, Capital Medical University, Beijing, China National Clinical Research Center for Neurological Diseases, Beijing, and Advanced Innovation Center for Human Brain Protection, Beijing Tiantan Hospital, Capital Medical University, Beijing, China (Y.W.).

Acknowledgment: The authors thank all study participants, their relatives, and the members of the survey teams at the 202 centers of the CHANCE-2 study.

Financial Support: By unrestricted grants from the Ministry of Science and Technology of the People's Republic of China, the Beijing Municipal Science and Technology Commission, the Chinese Stroke Association, the National Science and Technology Major Project (2017ZX09304018), and the Beijing Municipal Administration of Hospitals Incubating Program (PX2020021). Salubris Pharmaceuticals contributed ticagrelor, clopidogrel, and its placebo at no cost and with no restrictions. Chongqing Jingyin Bioscience provided the GMEX point-of-care genotyping system

and technical supports for CHANCE-2 at no cost and with no restrictions.

Disclosures: Disclosures can be viewed at www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum=M22-1667.

Data Sharing Statement: Data are available to researchers on request for the purposes of reproducing the results or replicating the procedure by directly contacting the corresponding authors.

Corresponding Authors: Xia Meng, MD, No. 119, South 4th Ring West Road, Fengtai District, Beijing 100070, China (e-mail, mengxia45@163.com); and Yongjun Wang, MD, No. 119, South 4th Ring West Road, Fengtai District, Beijing 100070, China (e-mail, yongjunwang@ncrcnd.org.cn).

Correction: This article was corrected on 1 November 2022 to correct a typographical error in the Methods section.

Author contributions are available at Annals.org.

References

1. Yalohom G, Schwartz R, Schwammenthal Y, et al. Chronic kidney disease and clinical outcome in patients with acute stroke. *Stroke*. 2009;40:1296-303. [PMID: 19182072] doi:10.1161/STROKEAHA.108.520882
2. Mostofsky E, Wellenius GA, Noheria A, et al. Renal function predicts survival in patients with acute ischemic stroke. *Cerebrovasc Dis*. 2009;28:88-94. [PMID: 19468220] doi:10.1159/000219302
3. Amarenco P, Callahan A 3rd, Campese VM, et al. Effect of high-dose atorvastatin on renal function in subjects with stroke or transient ischemic attack in the SPARCL trial. *Stroke*. 2014;45:2974-82. [PMID: 25147328] doi:10.1161/STROKEAHA.114.005832
4. Capodanno D, Angiolillo DJ. Antithrombotic therapy in patients with chronic kidney disease. *Circulation*. 2012;125:2649-61. [PMID: 22644369] doi:10.1161/CIRCULATIONAHA.111.084996

5. Bath PM, Woodhouse LJ, Appleton JP, et al; TARDIS Investigators. Antiplatelet therapy with aspirin, clopidogrel, and dipyridamole versus clopidogrel alone or aspirin and dipyridamole in patients with acute cerebral ischaemia (TARDIS): a randomised, open-label, phase 3 superiority trial. *Lancet*. 2018;391:850-9. [PMID: 29274727] doi:10.1016/S0140-6736(17)32849-0
6. Wang Y, Wang Y, Zhao X, et al; CHANCE Investigators. Clopidogrel with aspirin in acute minor stroke or transient ischemic attack. *N Engl J Med*. 2013;369:11-9. [PMID: 23803136] doi:10.1056/NEJMoa1215340
7. Johnston SC, Easton JD, Farrant M, et al; Clinical Research Collaboration, Neurological Emergencies Treatment Trials Network, and the POINT Investigators. Clopidogrel and aspirin in acute ischemic stroke and high-risk TIA. *N Engl J Med*. 2018;379:215-25. [PMID: 29766750] doi:10.1056/NEJMoa1800410
8. Kernan WN, Ovbiagele B, Black HR, et al; American Heart Association Stroke Council, Council on Cardiovascular and Stroke Nursing, Council on Clinical Cardiology, and Council on Peripheral Vascular Disease. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45:2160-236. [PMID: 24788967] doi:10.1161/STR.0000000000000024
9. Husted S, Emanuelsson H, Heptinstall S, et al. Pharmacodynamics, pharmacokinetics, and safety of the oral reversible P2Y12 antagonist AZD6140 with aspirin in patients with atherosclerosis: a double-blind comparison to clopidogrel with aspirin. *Eur Heart J*. 2006;27:1038-47. [PMID: 16476694] doi:10.1093/eurheartj/ehi754
10. Storey RF, Husted S, Harrington RA, et al. Inhibition of platelet aggregation by AZD6140, a reversible oral P2Y12 receptor antagonist, compared with clopidogrel in patients with acute coronary syndromes. *J Am Coll Cardiol*. 2007;50:1852-6. [PMID: 17980251] doi:10.1016/j.jacc.2007.07.058
11. Johnston SC, Amarenco P, Denison H, et al; THALES Investigators. Ticagrelor and aspirin or aspirin alone in acute ischemic stroke or TIA. *N Engl J Med*. 2020;383:207-17. [PMID: 32668111] doi:10.1056/NEJMoa1916870
12. Wang Y, Meng X, Wang A, et al; CHANCE-2 Investigators. Ticagrelor versus clopidogrel in CYP2C19 loss-of-function carriers with stroke or TIA. *N Engl J Med*. 2021;385:2520-30. [PMID: 34708996] doi:10.1056/NEJMoa2111749
13. McDermott JH, Leach M, Sen D, et al. The role of CYP2C19 genotyping to guide antiplatelet therapy following ischemic stroke or transient ischemic attack. *Expert Rev Clin Pharmacol*. 2022;15:811-25. [PMID: 35912831] doi:10.1080/17512433.2022.2108401
14. Best PJ, Steinhilber SR, Berger PB, et al; CREDO Investigators. The efficacy and safety of short- and long-term dual antiplatelet therapy in patients with mild or moderate chronic kidney disease: results from the Clopidogrel for the Reduction of Events During Observation (CREDO) trial. *Am Heart J*. 2008;155:687-93. [PMID: 18371477] doi:10.1016/j.ahj.2007.10.046
15. Zhou Y, Pan Y, Wu Y, et al; CHANCE Investigators. Effect of estimated glomerular filtration rate decline on the efficacy and safety of clopidogrel with aspirin in minor stroke or transient ischemic attack: CHANCE substudy (clopidogrel in high-risk patients with acute non-disabling cerebrovascular events). *Stroke*. 2016;47:2791-6. [PMID: 27738237] doi:10.1161/STROKEAHA.116.014761
16. Keltai M, Tonelli M, Mann JF, et al; CURE Trial Investigators. Renal function and outcomes in acute coronary syndrome: impact of clopidogrel. *Eur J Cardiovasc Prev Rehabil*. 2007;14:312-8. [PMID: 17446813] doi:10.1097/01.hjr.0000220582.19516.a6
17. James S, Budaj A, Aylward P, et al. Ticagrelor versus clopidogrel in acute coronary syndromes in relation to renal function: results from the Platelet Inhibition and Patient Outcomes (PLATO) trial. *Circulation*. 2010;122:1056-67. [PMID: 20805430] doi:10.1161/CIRCULATIONAHA.109.933796
18. Wang Y, Johnston C, Bath PM, et al; CHANCE-2 Investigators. Clopidogrel with aspirin in High-risk patients with Acute Non-disabling Cerebrovascular Events II (CHANCE-2): rationale and design of a multicentre randomised trial. *Stroke Vasc Neurol*. 2021;6:280-5. [PMID: 33952670] doi:10.1136/svn-2020-000791
19. Levey AS, Stevens LA, Schmid CH, et al; CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration). A new equation to estimate glomerular filtration rate. *Ann Intern Med*. 2009;150:604-12. [PMID: 19414839] doi:10.7326/0003-4819-150-9-200905050-00006
20. Teo BW, Xu H, Wang D, et al. GFR estimating equations in a multiethnic Asian population. *Am J Kidney Dis*. 2011;58:56-63. [PMID: 21601325] doi:10.1053/j.ajkd.2011.02.393
21. Levey AS, Coresh J, Balk E, et al; National Kidney Foundation. National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Ann Intern Med*. 2003;139:137-47. [PMID: 12859163] doi:10.7326/0003-4819-139-2-200307150-00013
22. Stevens PE, Levin A; Kidney Disease: Improving Global Outcomes Chronic Kidney Disease Guideline Development Work Group Members. Evaluation and management of chronic kidney disease: synopsis of the kidney disease: improving global outcomes 2012 clinical practice guideline. *Ann Intern Med*. 2013;158:825-30. [PMID: 23732715] doi:10.7326/0003-4819-158-11-201306040-00007
23. GUSTO Investigators. An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. *N Engl J Med*. 1993;329:673-82. [PMID: 8204123] doi:10.1056/NEJM199309023291001
24. Newcombe RG. Interval estimation for the difference between independent proportions: comparison of eleven methods. *Stat Med*. 1998;17:873-90. [PMID: 9595617] doi:10.1002/(sici)1097-0258(19980430)17:8<873::aid-sim779>3.0.co;2-i
25. Kim SJ, Bang OY. Antiplatelet therapy for preventing stroke in patients with chronic kidney disease. *Contrib Nephrol*. 2013;179:119-29. [PMID: 23652455] doi:10.1159/000346730
26. McClellan W, Aronoff SL, Bolton WK, et al. The prevalence of anemia in patients with chronic kidney disease. *Curr Med Res Opin*. 2004;20:1501-10. [PMID: 15383200] doi:10.1185/030079904X2763
27. Grassi G, Quarti-Trevano F, Seravalle F, et al. Early sympathetic activation in the initial clinical stages of chronic renal failure. *Hypertension*. 2011;57:846-51. [PMID: 21300663] doi:10.1161/HYPERTENSIONAHA.110.164780
28. Hansson EC, Jidéus L, Åberg B, et al. Coronary artery bypass grafting-related bleeding complications in patients treated with ticagrelor or clopidogrel: a nationwide study. *Eur Heart J*. 2016;37:189-97. [PMID: 26330426] doi:10.1093/eurheartj/ehv381
29. Pulcinelli FM, Pignatelli P, Celestini A, et al. Inhibition of platelet aggregation by aspirin progressively decreases in long-term treated patients. *J Am Coll Cardiol*. 2004;43:979-84. [PMID: 15028353] doi:10.1016/j.jacc.2003.08.062
30. Wallentin L, Becker RC, Budaj A, et al; PLATO Investigators. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med*. 2009;361:1045-57. [PMID: 19717846] doi:10.1056/NEJMoa0904327
31. Gurbel PA, Bliden KP, Butler K, et al. Randomized double-blind assessment of the ONSET and OFFSET of the antiplatelet effects of ticagrelor versus clopidogrel in patients with stable coronary artery disease: the ONSET/OFFSET study. *Circulation*. 2009;120:2577-85. [PMID: 19923168] doi:10.1161/CIRCULATIONAHA.109.912550
32. Teng R, Oliver S, Hayes MA, et al. Absorption, distribution, metabolism, and excretion of ticagrelor in healthy subjects. *Drug Metab Dispos*. 2010;38:1514-21. [PMID: 20551239] doi:10.1124/dmd.110.032250
33. Jiang XL, Samant S, Lewis JP, et al. Development of a physiology-directed population pharmacokinetic and pharmacodynamic

- model for characterizing the impact of genetic and demographic factors on clopidogrel response in healthy adults. *Eur J Pharm Sci.* 2016;82:64-78. [PMID: 26524713] doi:10.1016/j.ejps.2015.10.024
34. **Stefanini GG, Briguori C, Cao D, et al.** Ticagrelor monotherapy in patients with chronic kidney disease undergoing percutaneous coronary intervention: TWILIGHT-CKD. *Eur Heart J.* 2021;42:4683-93. [PMID: 34423374] doi:10.1093/eurheartj/ehab533
35. **Herzog CA, Shroff GR.** Atherosclerotic versus nonatherosclerotic evaluation: the Yin and Yang of cardiovascular imaging in advanced chronic kidney disease [Editorial]. *JACC Cardiovasc Imaging.* 2014;7:729-32. [PMID: 25034922] doi:10.1016/j.jcmg.2013.12.016
36. **Ocak G, Rookmaaker MB, Algra A, et al; SMART Study Group.** Chronic kidney disease and bleeding risk in patients at high cardiovascular risk: a cohort study. *J Thromb Haemost.* 2018;16:65-73. [PMID: 29125709] doi:10.1111/jth.13904
37. **Bonello L, Angiolillo DJ, Aradi D, et al.** P2Y12-ADP receptor blockade in chronic kidney disease patients with acute coronary syndromes. *Circulation.* 2018;138:1582-96. [PMID: 30354508] doi:10.1161/CIRCULATIONAHA.118.032078
38. **VanderWeele TJ, Knol MJ.** Interpretation of subgroup analyses in randomized trials: heterogeneity versus secondary interventions. *Ann Intern Med.* 2011;154:680-3. [PMID: 21576536] doi:10.7326/0003-4819-154-10-201105170-00008

Author Contributions: Conception and design: A. Wang, X. Xie, S.C. Johnston, H. Li, J. Jing, Yilong Wang, X. Zhao, Y. Jiang, X. Meng, Yongjun Wang.

Analysis and interpretation of the data: A. Wang, X. Xie, X. Tian, S.C. Johnston, P.M. Bath, Y. Zuo, Y. Jiang, L. Liu.

Drafting of the article: A. Wang, X. Xie, X. Tian, L. Liu.

Critical revision of the article for important intellectual content: A. Wang, S.C. Johnston, H. Li, P.M. Bath, X. Zhao, Z. Li, Y. Jiang, Yongjun Wang.

Final approval of the article: A. Wang, X. Xie, X. Tian, S.C. Johnston, H. Li, P.M. Bath, Y. Zuo, J. Jing, J. Lin, Yilong Wang, X. Zhao, Z. Li, Y. Jiang, L. Liu, X. Meng, Yongjun Wang.

Provision of study materials or patients: J. Lin, X. Meng, Yongjun Wang.

Statistical expertise: A. Wang, Y. Jiang.

Obtaining of funding: X. Zhao, X. Meng, Yongjun Wang.

Administrative, technical, or logistic support: Yilong Wang, Y. Jiang, X. Meng, Yongjun Wang.

Collection and assembly of data: A. Wang, X. Xie, J. Jing, J. Lin, Y. Jiang, X. Meng.